Searching PAJ Page 1 of 1

## PATENT ABSTRACTS OF JAPAN

(11)Publication number:

05-194192

(43) Date of publication of application: 03.08.1993

(51)Int.Cl.

A61K 9/127 A61K 47/24 B01J 13/02

(21)Application number : 04-215550 (22)Date of filing:

20.07.1992

(71)Applicant: SHIONOGI & CO LTD

(72)Inventor: JIZOUMOTO HIROAKI

HIRANO KOICHIRO KANEOKA ERI

(30)Priority

Priority number: 40323541

Priority date: 21.08.1991 Priority country: JP

### (54) TEMPERATURE-SENSITIVE MLV-TYPE LIPOSOME WITH IMPROVED DISPERSIBILITY

(57)Abstract:

PURPOSE: To provide the subject liposome excellent in the controlled releasability for medicine, useful for cancer, tumor, etc., comprising dipalmitoylphosphatidyl- choline and negatively charged diacyl-glycerophosphatide at a specified proportion.

CONSTITUTION: The objective liposome comprising (A) dipalmitoylphosphatidyl- choline and (B) a negatively charged 1,2-diacyl-glycerophosphatide (where, acyl groups at 1- and 2-sites are each palmitoyl or stearoyl independent of each other) at the weight ratio of (20:1) to (8:2). This liposome has the following characteristics: (1) the osmotic pressure ratio for inside aqueous phase is 1.5 to 2 times that for physiological isotonic liquid, and (2) particle diameter is 0.03-0.5μm.

#### \* NOTICES \*

JPO and INPIT are not responsible for any damages caused by the use of this translation.

- 1. This document has been translated by computer. So the translation may not reflect the original precisely.
- 2.\*\*\*\* shows the word which can not be translated.
- 3.In the drawings, any words are not translated.

#### CLAIMS


#### [Claim(s)]

[Claim 1]A 1,2-diacyl glycerophospholipid which has dipalmitoylphosphatidylcholine and a negative charge (however, acyl of the 1st place and the 2nd place) mutual -- independently -palmitoyl or stearoyl -- it is -- the temperature sensitivity MLV type liposome by which dispersibility containing by a weight ratio of 20:1-8:2 has been improved.

IClaim 2|The temperature sensitivity MLV type liposome according to claim 1 whose osmotic pressure ratio of inner aqueous phase is twice [ 1.5 to ] the physiological isotonic solution. [Claim 3]The temperature sensitivity MLV type liposome according to claim 1 or 2 whose particle diameter is 0.03-0.5 micrometer.

[Translation done.]

# Drawing selection Drawing 1 -



[Translation done.]